

Intrauterine bone findings during hysteroscopy at GESHRTH – Yaoundé: report of six cases

Trouvailles endocavitaires osseuses durant l'hystérocopie au CHRACERH : à propos de six cas

Nyada SR^{1,2}, Mpono P^{1,2}, Ngono Akam V^{1,2}, Ngassam A^{1,2}, Kasia Onana YB¹, Mendibi S¹, Belinga E^{1,2}, Noa Ndoua CC^{1,2}, Kasia JM^{1,2}

Original Article

1. Gynecological Endoscopic Surgery and Human Reproduction Teaching Hospital of Yaoundé (GESHRTHY)
2. Faculty of Medicine and Biomedical Sciences – University of Yaoundé 1

Auteur correspondant :

Nyada Serge Robert, tel (00237) 677600154. Email : sergenyadar@yahoo.fr

Key words: Bone, Hysteroscopy, GESHRTHY, Cameroon

Mots-clés : Séquestres osseux, Hystérocopie, CHRACERH, Cameroun.

Date de soumission: 04/02/2026

Date d'acceptation: 06/04/2026

ABSTRACT

Introduction: The presence of intrauterine bone structures during hysteroscopy is an unusual finding.

Materials and Methods: We carried out a retrospective descriptive study at the Gynecological Endoscopic Surgery and Human Reproduction Teaching Hospital of Yaoundé (GESHRTHY) between september 1st 2017 to august 31st 2022. A public facility addressing infertility and dedicated to minimally invasive surgery in Yaoundé. All procedures of hysteroscopy performed in the theatre were reviewed. All cases of patients reporting the presence of intrauterine bones during the procedure were processed.

Results: We found six cases of intrauterine bone structures out of 854 hysteroscopies. The prevalence over the 5-year period was 0,7%. The mean age of patients was 35.2 years. All patients revealed a past history of mid trimester abortion. For one patient, the history of abortion was confessed retrospectively after the diagnosis of retained bony fragments. A past history of Sexually Transmitted Infection was recorded for three patients (50%). The clinical presentation of patients included infertility in all patients (100%), lower abdominal pain in three (50%) and oligomenorrhoea in two (33.3%). Routine work-up identified the presence of intra uterine calcifications in five patients (83.3%). In one case, the finding of bone structure was fortuitous. The hysteroscopic procedure permitted the complete retrieval of all bone fragments. There were no complications.

Conclusion: The prevalence of intra uterine bone findings during hysteroscopy was 0.7%. All patients complained of infertility and recorded a past history of mid trimester abortion.

RESUME

Introduction : La présence intra utérine de fragments osseux au cours d'une hystérocopie constitue une trouvaille inhabituelle.

Méthodologie : Nous avons mené une étude descriptive rétrospective au Centre Hospitalier de Recherche et d'Application en Chirurgie Endoscopique et Reproduction Humaine de Yaoundé (CHRACERH) entre le 1er septembre 2017 et le 31 août 2022. Nous avons revu tous les cas d'hystérocopie et enrôlé toutes les patientes chez qui la présence de débris osseux intra cavitaires était rapportée.

Résultats : Nous avons retenu six cas parmi 854 procédures. La fréquence de séquestres osseux durant la période d'étude de 5 ans était de 0,7%. L'âge moyen des patientes étaient de 35,2 ans. Toutes les patientes avaient un antécédent d'avortement tardif. Pour une patiente, l'antécédent d'avortement n'était avoué qu'après la chirurgie et au vu des trouvailles opératoires. Un antécédent d'infection sexuellement transmissible était retrouvé chez trois patientes (50%). La présentation clinique des patientes comportait une infertilité dans 100%, une pelvi-algie dans 50% et une oligoménorrhée dans 33% des cas. Les examens de routine ont identifié la présence de calcifications intra-utérines dans 5 cas (83,3%). Chez une patiente, la découverte de séquestres osseux était fortuite. Nous avons au cours de l'hystérocopie procédé à l'ablation complète de tous les fragments osseux. Nous n'avons enregistré aucune complication.

Conclusion : La fréquence des fragments osseux de découverte hystérocopique était de 0,7%. Toutes les patientes se plaignaient d'infertilité et avaient un antécédent d'avortement tardif. L'hystérocopie a permis l'ablation complète des séquestres.

DOI : <https://doi.org/10.64294/jsd.v4i2.332>

Introduction

The presence of foreign bodies such as bone structures within the uterine cavity is an uncommon hysteroscopic finding. The incidence of bone retention among women undergoing diagnostic hysteroscopy is 0.15% [1]. Bone fragments account to 11.9% of foreign bodies removed from uterus by hysteroscopy of infertile patients [2]. In the literature, there are two types of bone structures within the uterine cavity: the retention of foetal bones and the osseous metaplasia of endometrium.

Many theories can explain the presence of intra uterine bones structures. Foetal bone retention is described as the uterine retention of foetal bones secondary to an incomplete second trimester abortion. Endometrial osseous metaplasia, defined as the presence of mature or immature bone tissue within the endometrium can be explained by several theories [3]. The most widely accepted theory is endochondral ossification from omnipotent endometrial cells, derived from embryonic remnants. Another theory is related to stromal mesenchymal cells graft from foetal origin, which can differentiate into osteoblasts after termination of pregnancy, via cytokines and growth factors. The last theory which has been described in women who have never been pregnant is true osseous metaplasia of endometrial cells into osteoblast cells in response to chronic inflammation or trauma, similar to that occurring after calcification of fibrosis or abscesses [3-7].

Patients usually have no specific symptoms. Case reports have highlighted infertility, abnormal uterine bleeding, lower abdominal pain, dyspareunia, dysmenorrhoea, vaginal discharge and spontaneous passage of bone fragments in menstrual blood as major symptoms [2,7]. The main objective of this study was to determine the frequency and clinical presentation of intrauterine bone findings among women attending hysteroscopy in our settings.

Materials and Methods

A retrospective descriptive study was carried out at the GESHRTHY between september 1st 2017 to august 31st 2022. The files of all women who underwent diagnostic or operative hysteroscopy were reviewed. The patients presenting intrauterine bony findings during the procedure were identified. Data from medical files and reports were collected, including past medical history, presenting symptoms, pelvic ultrasound and radiological findings, surgical procedure data and postoperative follow up. Histopathological analysis of specimen collected during the surgery was requested. Written signed consent was obtained from all patients before the surgery. Confidentiality was guaranteed during the whole procedure. Approval from the ethic committee board was not sought because of the descriptive nature of the study and no patient was subjected to

any other interventions.

Results

During the period of the study, 854 hysteroscopies were performed. Among these, six (6) cases of intrauterine bony retention were identified. Thus, the frequency was 0.7%. The median age of patients was 35.2 years (range: 28 - 40 years). All patients had a past history of mid trimester abortion with surgical approach dating between three to 16 years. Two patients (33.3%) had a history of three previous mid trimester abortion. However, in one case, the history of abortion was confessed after the surgical procedure and in view of the operative findings.

A past history of Sexually Transmitted Infection (STI) was recorded in three cases (50%). It was mycoplasma infection in two cases (33.3%) and chlamydial infection in one case (16.6%). On clinical presentation (Fig 1), all patients complained of infertility; the average duration was 6.3 years (range 2-12 years). Chronic lower abdominal pain was identified in three cases (50%) and oligomenorrhoea in two cases (33.3%).

The presence of endometrial calcifications was recorded in four cases (66.7%) out of six pelvic ultrasounds, in two cases (66.7%) out of three hysterosonographies and in one case (50%) out of two hysterosalpingographies, alongside with nonspecific findings such as fibroids and polyps (Fig 2). It should be noted that in one case, no intrauterine abnormality other than calcified intramural fibroids was found on routine scanning.




Fig 1: clinical presentation (N=6)




Fig 2: routine scanning findings

Complete hysteroscopic retrieval of bony fragments was achieved in all cases (Fig 3-5). However, one patient needed two procedures to reach that goal. The average number of bone fragments was 8 (range 6-20). These were recognizable long and flat bones

with nearly intact morphology. In one case however, a coral-like bone was described. During the procedure as well, stage 1 synechia was observed and treated in two cases (33.3%).

The postoperative follow-up was unremarkable. All patients resumed menses with normal volume during the next menstrual cycle and the ultrasound performed in all patients after one month was normal. Pathological analysis of the endometrial specimen was carried out in three patients and reported a chronic endometritis with calcification. No DNA analysis was done.

diagnosed between eight weeks to 19 years after the termination of pregnancy [2,9]. Our study suggest that all patients had a past history of mid trimester abortion with surgical approach dating between three to 16 years and two patients (33.3%) had a history of three previous mid trimester abortions. One patient did not even mention the history of abortion until after the hysteroscopy was performed. The reason being the legal settings, as in some developing countries, abortion is prohibited. This leads to almost all abortions being illegally performed. There should be a number of undiagnosed women with this condition in the community. Therefore, it is expected a higher incidence of intrauterine retained bone fragments could be revealed by improving hysteroscopic practice. Physicians should keep in mind the remote possibility of a forgotten abortion while assessing infertility in our settings.

Xiao et al. emphasized on the need of a systematic B-ultrasound examination to ensure the emptiness of the uterus in case of termination of a pregnancy older than 12 weeks [10]. Gainder et al. found that the main symptoms were menorrhagia (five cases), hypo/amenorrhoea (two cases), metrorrhagia (four cases) and chronic pelvic pain (two cases). They reported an incidental diagnosis of foetal bone in 12/18 patients [8]. Van Den Bosch et al. suggested that the presence of intramural bone does not impair fertility [11]. Makris et al reported irregular vaginal bleeding as main complain of patients in his study [1]. Recurrent vaginitis and pelvic inflammatory disease may be rare presentations of foetal bone retention [2]. Our study outlined infertility, lower abdominal pain and oligomenorrhoea as part of clinical symptoms in 100%, 50% and 33% respectively. The fact that our facility is dedicated to infertility can explain that all our patients complained of infertility.

Retained foetal bones are supposed to act the same way as an intrauterine contraceptive by increasing the production of menstrual prostaglandin E and prostacyclin. Increased level of prostaglandin E may also be associated with pelvic pain and dysmenorrhoea [2,9]. It has been demonstrated that prostaglandin E2 concentrations decreased by 50% after removal of bone [9].

The diagnosis is often unexpected on routine ultrasound. On ultrasonographic routine examination, the presence of intrauterine bright echogenic area with posterior acoustic shadowing should alert the physician [6]. However, differential diagnoses include intrauterine devices, foreign bodies, Asherman's syndrome, calcified submucosal fibroids, mixed mesodermal tumours and adenosarcoma [2,9]. The report of the ultrasound scan of one of our patients related the sole presence of calcified intramural fibroids. We think that the ultra-sonographer did not pay attention to the presence of intrauterine




Fig 3: hysteroscopic view of bones: A and B (Arrow shows bones)




Fig 4: Hysteroscopic view of bones: (Arrow shows bones)
A: Hysteroscopic view
B: Hysteroscopic retrieval of bone with grasper




Fig 5: Post operative view of the bones retrieved: A and B

Discussion

The incidence rates of retained intrauterine foetal bones found by Makris et al. and Gainder et al. were respectively 0.15% and 0.28% [1,8]. A higher incidence of 0.7% was revealed in our actual study. Gainder et al. reported a past history of first and second trimester abortion in five and 13 cases respectively out of 18 [8]. In the literature, foetal bone retention has been

calcifications allusive of bone retention due to the presence of numerous intramural calcified fibroids.

For definitive treatment, hysteroscopy is the gold standard. It may be helped by a preoperative gonadotropin-releasing hormone analogues therapy, which permit to see all the abnormal tissue hence facilitate the removal [1,2]. The use of ultrasound is advocated in case of extensive metaplasia or numerous fragments to ensure safety and complete removal of bone fragments [9,10].

The presence of bone marrow in the fragments is believed to be an argument against foetal origin since bone is not populated with hematopoietic cells until the third trimester of pregnancy. Although a heterotopic bone might be colonized by circulating hematopoietic stem cells leading to mature marrow. This could be definitely resolved by DNA analysis of the fragments [7]. In the literature, many case reports suggested the solving of infertility issue after successful treatment of intrauterine bone retention [10,12].

Conclusion

The prevalence of intrauterine bone fragments during hysteroscopy at GESHRTHY was 0.7% over a 5-years period. Infertility and a past history of mid trimester abortion were recorded in all cases. Routine scanning allowed the diagnosis of intrauterine calcifications in 83.3%. Hysteroscopy permitted a successful and safe retrieval of all bone fragments.

Interest disclosure: The authors declare that there is no conflict of interest regarding the publication of this paper.

Author's contribution: Nyada Serge: design, data acquisition, analysis, manuscript write up and edition. All the authors have read and agreed to the final manuscript.

References

- Makris N, Stefanidis K, Loutradis D, Anastasiadou K, Hatjipappas G, Antsaklis A. The incidence of retained fetal bone revealed in 2000 diagnostic hysteroscopies. *JLS*. 2006; 10: 76-77.
- Demirtas O, Terzi H, Kale A, Sanibrahim B, Tolga Guler O. A rare and unexpected case of retained fetal bone after an unsafe abortion. *J Pak Med Assoc*. 2015; 65 (10) :1119 -1121
- Saponara S, Angioni S, Parry J P, Pacheco L A, Carugno J, Moawad N, et al. The pivotal role of hysteroscopy in diagnosing subtle uterine lesions in infertile patients: Seeing the unseen can make the difference, *Eur J Obstet Gynecol Reprod Biol*. 2025; 305: 132-141. doi: <https://doi.org/10.1016/j.ejogrb.2024.12.013>
- Fernandez H, Garbin O, Gervaise A. *Hystérocopie et fertiiloscopie*. Elsevier Masson, Techniques chirurgicales – Gynécologie. 2013 : 208-209.
- Connor ME, Clark TJ. *Diagnostic and operative hysteroscopy*. Cambridge University Press. 2020; 51-53
- Sardo ADS, Guida M, Bettocchi S, Nappi L, Sorrentino F, Bifulco G, Nappi C. Role of hysteroscopy in evaluating chronic pelvic pain. *Fertil Steril*. 2008 ; 90(4) :1191–6. DOI: 10.1016/j.fertnstert.2007.07.1351
- Basu M, Mammen C, Owen E. Bony fragments in the uterus: an association with secondary subfertility. *Ultrasound Obstet Gynecol* 2003; 22:402-406. DOI: 10.1002/uog.221
- Gainder S, Arora P, Dhaliwal LK. Retained intrauterine bony fragments as a cause of secondary infertility in a tertiary level Indian hospital. *J Hum Reprod Sci* 2018; 11:286-290.
- Gulec UK, Parlakgumus HA, Kiliçdag EB, Bolat F, Bagis T. Osseous metaplasia of the endometrium. *BMJ Case Rep*. 2010;10.1136/bcr.04.2010.2931
- Xiao S, Tian Q, Xue M. Infertility caused by intrauterine fetal bone retention: a case report. *J Med Case Rep*. 2014; 8: 177. DOI:10.1186/1752-1947-8-177
- Van Den Bosch T, Van Schoubroeck D, Timmerman D, Deprest J. Uterine intramural bone after mid-trimester termination of pregnancy may not affect fertility: a case report. *Ultrasound Obstet Gynecol*. 2003; 22: 407–408. DOI: 10.1002/uog.220
- Mishra N, Bharti R, Mittal P, Suri J, Pandey D. Retained intra-uterine foetal bones resulting in secondary infertility: a case report. *Cureus*. 2018; 10(5): e2575. DOI 10.7759/cureus.2575